

Influence of Fetal Origin of Posterior Cerebral Artery on Ischemic Stroke

VIGNESHWAR ADHITHIYA¹, KULASEKARAN NADHAMUNI², MOHAMED RAFI KATHAR HUSSAIN³, AM ANAND⁴, UMAMAGESWARI AMIRTHALINGAM⁵, ELAMPARIDHI PADMANABAN⁶



ABSTRACT

Introduction: Fetal origin of posterior cerebral artery is one among the common variants in the Circle of Willis. In case of cerebral injury due to thrombo-embolic stroke, the presence of fetal type of PCA may modify its severity and the territory involved. The purpose of this study is to review and to evaluate the potential clinical significance of fetal type PCA in ischemic stroke.

Aim: To analyse whether fetal PCA and its types are associated with ischemic stroke.

Materials and Methods: This is an analytical cross-sectional study conducted from January 2019 to January 2020 to assess the PCA in the 174 patients referred to Magnetic Resonance Imaging

(MRI) Brain and diagnosed as ischemic stroke. Data was entered using excel sheet and analysed using R v.3. 5.1. Descriptive analysis was performed followed by bivariate analysis.

Results: There is no statistically significant relationship between the presence of fetal PCA and ischemic stroke and the p-value is 0.936.

Conclusion: There is no significant association of presence or absence of partial or complete type of fetal PCA with ischemic stroke. No significant association was also found between the presence of fetal PCA and vascular territory of infarct and laterality of ischemic stroke.

Keywords: Anatomical variants, Circle of willis, Infarct, MRA

INTRODUCTION

The circle of Willis has many variants, one of the common variants being fetal origin of the Posterior Cerebral Artery (PCA) from the Internal Carotid Artery (ICA) [1]. Angiographic studies reveals that 11% to 46% of adults will have fetal type of PCA either unilaterally or bilaterally [2]. Fetal type PCA is defined as a PCA that originates from the ICA with or without a small connection with the Basilar Artery (BA). In this condition, instead of the BA, the ICA supplies blood to the PCA [3]. Collateral circulation in the brain is important for maintaining a sufficient cerebral blood flow in case of ischemic stroke [1].

In case of cerebral injury due to thromboembolic stroke, the presence of fetal type of PCA may modify its severity and the territory involved [3].

MATERIALS AND METHODS

The present analytical cross-sectional study was conducted at Sri ManakulaVinayagar Medical College and hospital, Puduchery, which is a multidisciplinary, 900 bedded hospital with fully equipped Radio-diagnosis department. The study was conducted during the period of January 2019 to January 2020, to assess the posterior cerebral arteries in the 174 patients referred to MRI Brain. The inclusion criteria used in this study were those patients with diagnosis of ischemic stroke on MRI. Patients with history of head trauma, cerebral surgery, vascular malformations, vasculitis and bilateral fetal PCA were excluded. The MRI images were acquired in the axial, coronal and sagittal planes using 1.5 Tesla PHILIPS whole body MR systems with standard imaging head coil with standard stroke protocol. Magnetic Resonance Angiography (MRA) was also done to assess the presence and type of fetal PCA. The primary outcome of this study was to evaluate the potential clinical significance of fetal type PCA in ischemic stroke by comparing with normal type PCA.

STATISTICAL ANALYSIS

Data was entered using excel and analysed using R v.3. 5.1. Descriptive analysis was performed followed by bivariate analysis.

The p-value was calculated using single proportion test, chi-square test or Fisher-exact test as appropriate. The p-value less than 0.05 was considered statistically significant.

RESULTS

Among the 174 cases studied, 32 (18.4%) patients were below 40 years of age, 74 (42.5 %) patients were above the age of 60 years and 68 (39.1 %) patients were between the age of 41 to 60. Of the 174 MRI studies reviewed, it was found that 68 (39.1%) patients were having fetal PCA compared to the remaining 106 (60.9 %) who had normal posterior circulation. In this study, male patients were more (66.7%) when compared to female patients. Among the 116 males, 48 (41.3%) were found to have fetal PCA and among 58 females, 20 (34.4%) were found to have fetal PCA. Among the patients with fetal PCA, 38 (55.9%) patients had partial type of fetal PCA and 30 (44 %) patients had complete type of fetal PCA.


In this study, when comparing the side of fetal type of PCA, the right side fetal PCA is seen in 40 (58.8%) patients, which was more common than the left side fetal PCA which is seen in 28 (41.2%) patients.

Of the 174 patients, 140 (80.5%) patients had Middle Cerebral Artery (MCA) territory infarct and 34 (19.5%) patients had PCA territory infarct.

The distribution of type of PCA with respect to vascular territory of the infarct is given in [Table/Fig-1]. There was no significant association between the presence of fetal PCA and vascular territory of infarct (p-value is 0.936).

Among the patients with fetal PCA, ipsilateral infarcts were seen in 42 (61.8%) patients and contralateral infarcts were seen in 26 (38.2%) patients. There was no significant association between the presence of fetal PCA with ipsilateral or contralateral ischemic stroke (p-value is 0.229).

The distribution of the type of Fetal PCA with respect to the laterality of infarct is shown in [Table/Fig-2]. There was no statistically



[Table/Fig-1]: Distribution of type of PCA with respect to vascular territory of the infarct (p-value is 0.936).
PCA: Posterior cerebral artery; MCA: Middle cerebral artery

	Ipsilateral side	Contralateral side	p-value
Partial type	22	16	0.867
Complete type	20	10	

[Table/Fig-2]: Distribution of the type of fetal PCA in accordance with laterality of infarct. Chi-square test applied

significant relationship between the type of fetal PCA and laterality of infarct and the p-value is 0.867.


DISCUSSION

The cerebral blood flow can be divided into two circulations: the anterior and the posterior circulation [4]. The posterior circulation comprises of vertebral arteries and its branches and BA which bifurcates into PCAs on each side. The PCA has three segments, named as P1, P2 and P3. P1 segment lies between the basilar bifurcation and the posterior communicating artery. The P2 segment lies between the posterior communicating artery and the posterior aspect of the midbrain and P3 segment of PCA starts from the pulvinar to the anterior limit of the calcarine fissure [5,6]. The term complete fetal type PCA is defined as the total absence of P1 segment of PCA, as shown in [Table/Fig-3]. Whereas, the term partial type fetal PCA is defined as hypoplastic P1 segment, as shown in [Table/Fig-4]. This P1 segment is the segment between basilar bifurcation and post communicating part of PCA [7,8].

In case of fetal PCA, the leptomeningeal vessels fails to develop between anterior and posterior circulation, hence there is more risk



[Table/Fig-3]: MR angiogram Maximum Intensity Projection (MIP) image shows right complete type of fetal PCA (yellow arrow). Another anatomical variation seen in this case is hypoplastic right A1 segment (white arrow).




[Table/Fig-4]: MR angiogram Maximum Intensity Projection (MIP) image shows left hypoplastic P1 segment (arrow) with prominent posterior communicating artery- Suggestive of left partial fetal PCA.

of developing ischemic stroke. In partial fetal PCA, there is lack of development of leptomeningeal vessels as both the anterior and the posterior circulations are derived from the ICA, and collaterals could not be formed due to tentorium which prevents it. But a complete type fetal PCA can adapt to altered haemodynamics efficiently by developing collaterals [1]. Complete type of fetal PCA can impact the anatomy of the cerebral circulation by forming collaterals. Thus, in complete type PCA more area is perfused by the anterior circulation as PCA is completely supplied by ICA through collaterals [2]. Van Raamt AF et al., found lack of development of leptomeningeal collaterals in the fetal configuration of Circle of Willis, however their study did not show any association of full FTP with ischemic strokes [7].

In this study, the male patients were about 66.7% when compared to study done by Roshan KA et al., in which 60.3% patients were male which is close to present observations [9]. A 26% of patients had fetal type PCA in Roshan KA et al., study while this study shows 39.1% of fetal PCA, which is higher [9]. In a study done by Roopashree R, 20% percent of patients had fetal type of PCA while this study shows a higher percentage [10]. A 21% of fetal PCA patients in this study have partial type of PCA while Roshan KA et al., study shows 10% which is less [9].


When compared to normal PCA, there is no statistical significant association between the presence of fetal type PCA and MCA territory ischemic stroke in this study [Table/Fig-5].



[Table/Fig-5]: MRI axial sections of FLAIR, ADC and DWI sequences showing acute left MCA territory infarct.

Brandt T et al., study shows that incidence of PCA infarct is about 5% to 10% in general population [11]. This study shows 19% of ischemic infarcts involving the PCA territory as shown in [Table/Fig-6] which is higher in the population of this study. Commonly, the atherosclerotic diseases of posterior circulation will lead to PCA territory infarct, while in case of presence of fetal PCA, atherosclerotic plaque or embolisation of common carotid artery causes occipital infarct [12]. Argyro M et al., in their study also concluded that in the presence of fetal PCA anatomical variant, since ICA is the dominant blood supplier, thrombosis or embolism of the ICA may cause infarction of the occipital lobe [13].

No significant association between presence of fetal PCA and ischemic stroke was noted in this study which is similar



[Table/Fig-6]: MRI axial sections of FLAIR, ADC and DWI sequences showing acute right PCA territory infarct.

to the study done by Amre N et al., [14]. Roshan KA et al., also states that there is no significant association of partial or complete type of fetal PCA with ischemic stroke and the results are similar to present study. Van Raamt AF et al., found lack of development of leptomeningeal collaterals in the fetal configuration of Circle of Willis, however did not show any association of complete fetal type PCA with ischemic strokes [7]. Lima FO et al., detected no clinical influence of the presence of fetal PCA on the outcome of ischemic stroke [15]. Shaban A et al., studied the frequency and implication of stroke ipsilateral to fetal type PCA [16]. It also suggested that the presence of fetal type PCA does not increase the risk of ischemic stroke. Various studies have been done on impact of fetal type PCA on ischemic stroke. Among these studies van Raamt AF et al., and Shaban A et al., found no association of presence of fetal PCA and ischemic stroke [7,16]. This study show similar results. We could not find any association between stroke and presence of fetal type PCA.

Limitation(s)

This study was done using 1.5T MRI which was available in the institution. 3T MRI/CT angiogram/Digital subtraction angiography may give more spatial resolution, information and better outcome.

CONCLUSION(S)

There was no significant association found between presence of fetal PCA and ischemic stroke, and also no significant association found between the presence of fetal PCA or its types with ipsilateral or contralateral ischemic stroke.

Acknowledgement

We thank Dr. Lena Charlette, Assistant professor, Department of Community medicine, Andaman and Nicobar Islands Institute of Medical Science, for her support in the statistical part of the study.

REFERENCES

- [1] Prasanna VE, Balakrishnan R, Ramadoss K, Gnanashanmugham G. Can internal carotid artery occlusion produce simultaneous anterior and posterior circulation stroke? *Indian J Vasc Endovasc Surg.* 2015;2:130-32.
- [2] Jongen JC, Franke CL, Soeterboek AA, Versteeg CW, Ramos LM, Van J. Blood supply of the posterior cerebral artery by the carotid system on angiograms. *J Neurol.* 2002;249:455-60.
- [3] Stephen BS, Frank W, Zhora O, Lionel B, Anedward CM. Fetal-type variants of the posterior cerebral artery and concurrent infarction in the major arterial territories of the cerebral hemisphere. *J Investig Med High Impact Case.* 2016;4(3).
- [4] Lynn R, Abou-Chebl A. *Cerebral arterial anatomy. Practical carotid artery stenting.* London: Springer; 2009: pp.63-74.
- [5] Haines DE. *Fundamental neuroscience.* Philadelphia 2002.
- [6] Tatu L, Moulin T, Bogousslavsky J, Duvernoy H. Arterial territories of the human brain: cerebral hemispheres. *Neurology.* 1998;50:1699-708.
- [7] Van raamt AF, Mali WP, Laar PJ, Graaf Y. The fetal variant of the circle of Willis and its influence on the cerebral collateral circulation. *Cerebrovasc Dis.* 2006;22:217-24.
- [8] Arjal RK, Zhu T, Zhou Y. The study of fetal type posterior cerebral circulation on multislice CT angiography and its influence on cerebral ischemic strokes. *Clin Imaging.* 2014;38:22-25.
- [9] Roshan KA, Tingting Z, Yicheng Z. The study of fetal type posterior cerebral circulation on multislice CT angiography. *Clinical Imaging.* 2014;38:221-25.
- [10] Roopashree R. An anatomical study on relationship between posterior cerebral artery and posterior communicating artery. *International Journal of Anatomy, Radiology and Surgery.* 2013;2(1):9-12.
- [11] Brandt T, Steinke W, Thie A, Pessin MS, Caplan LR. Posterior cerebral artery territory infarcts: clinical features, infarct topography, causes and outcome. *Cerebrovasc Dis.* 2000;10:170-82.
- [12] Fisher CM. The posterior cerebral artery syndrome. *Can J Neurol Sci.* 1986;13:232-39.
- [13] Argyro M, Nikolaos E, Olympia P, Christina K, Dimitrios F, Aristodimos M, Nikolaos K, et al. High resolution MR angiography of the posterior cerebral circulation: variants, incidence and clinical impact. *Open Journal of Radiology.* 2013;3:209-14.
- [14] Amre N, Jessica R, Ruland S. Ischemic posterior circulation stroke: a review of anatomy, clinical presentations, diagnosis, and current management. *Frontiers in neurology.* 2014;Pp 12.
- [15] Lima FO, Furie KL, Silva GS. The pattern of leptomeningeal collaterals on CT angiography is a strong predictor of long-term functional outcome in stroke patients with large vessel intracranial occlusion. *Stroke.* 2010;41(10):2316-22.
- [16] Shaban A, Albright KC, Boehme AK, Martin-Schild S. Circle of Willis Variants: Fetal PCA. *Stroke Research and Treatment.* 2013;p.1-6.

PARTICULARS OF CONTRIBUTORS:

1. Junior Resident, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
2. Professor and Head, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
3. Assistant Professor, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
4. Assistant Professor, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
5. Associate Professor, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
6. Assistant Professor, Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

AM Anand,
Department of Radiodiagnosis, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India.
E-mail: anandmrd@gmail.com

PLAGIARISM CHECKING METHODS: [\(Jain H et al.\)](#)

- Plagiarism X-checker: Jun 26, 2020
- Manual Googling: Jul 30, 2020
- iThenticate Software: Nov 02, 2020 (07%)

ETYMOLOGY: Author Origin

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Jun 25, 2020**
Date of Peer Review: **Aug 05, 2020**
Date of Acceptance: **Aug 19, 2020**
Date of Publishing: **Jan 01, 2021**