

Computed Tomographic Findings of Malignant Bowel Neoplasms: A Retrospective Study

DEVARA ANIL KASHI VISHNUVARDHAN¹, SONICA SHARMA², AVINASH VERNEKAR³, CHANDHANA MERUGU⁴

ABSTRACT

Introduction: Malignant neoplasms of bowel particularly colorectal carcinomas are third commonest cause of death all over world. Computed Tomography (CT) plays a very important role in evaluation of malignant bowel neoplasms. CT examination for gastrointestinal tract evaluates the intestinal lumen along with demonstration of intestinal wall, adjacent tissues and solid viscera.

Aim: To evaluate malignant neoplasms of small and large intestine and also to define the various CT findings in malignant bowel neoplasms, extra-intestinal involvement to stage intestinal neoplasms and to accurately characterise the tumour histologically as far as possible.

Materials and Methods: A retrospective descriptive study included 38 patients who were proven to have malignancy of bowel on biopsy, underwent CT abdomen in major public hospital in a period of from November 2014 to May 2017 and evaluated for various CT findings in malignant bowel neoplasms in terms of involved intestine, length of diseased intestinal segment, degree of intestinal wall thickening, extra-intestinal involvement to stage bowel neoplasms and to accurately characterise the tumour histologically as far as possible. Data were tabulated and presented as frequency (n) and percentage (%).

Results: In Small intestine, ileum is the commonest site of involvement seen in 5 cases (13%), whereas rectum 10 cases (26.3%) was the commonest site of involvement in large intestine. Short segment involvement of bowel is more common 27 cases (71.1%) than long segment involvement 11 cases (28.9%). Circumferential wall thickening is more common in 33 cases (86.8%) than eccentric wall thickening 5 cases (13.2%). Lymph nodal involvement was seen in 17 cases (44%) and metastasis to liver was also seen in 11 cases (28.9%). Local invasion with involvement of surrounding structures was seen in 4 cases (10.5%). Adenocarcinoma was the commonest histological subtype seen in the highest number of cases 23 cases (60.5%), followed by lymphoma in 9 (23.6%) of cases, mucinous adenocarcinoma in 4 (10.5%) cases and few cases of about 2 (5.2%) were Gastrointestinal Stromal Tumour (GIST).

Conclusion: Computed Tomography (CT) imaging findings like involved intestine, length of diseased intestinal segment, degree of intestinal wall thickening, extra intestinal involvement with respect to involvement of surrounding structures, lymph nodal involvement, distant metastasis along with histological type in most cases could be found, thus permitting tumour characterisation which can provide very useful guide in diagnosis of malignant neoplasms of small and large intestine.

Keywords: Computed tomography, Large intestine, Metastasis, Small intestine, Wall thickening

INTRODUCTION

Abdominal studies were originally confined to solid organs in retro peritoneum extending eventually into peritoneal cavity [1]. Bowel details were not referred owing to slow data acquisition and lack of definition. Major improvement in computing science as well as development of spiral technique has now led to large volume data acquisition [2-4]. Thus, in recent times CT has been shown to be an essential tool in radiologic evaluation of gastrointestinal tracts [5]. Barium study and endoscopy remain the method of choice in initial detection of most of alimentary tract disease. However, both techniques are limited to examination of mucosal surface, caliber and contour of bowel segment. They provide little and indirect information regarding intramural or extrinsic abnormalities. CT examination for gastrointestinal tract not only evaluates the bowel lumen, but its major contribution consists of demonstrating bowel wall and the adjacent tissues and organs [6,7]. Malignant neoplasms of bowel particularly colorectal carcinomas are third commonest cause of death all over world. CT plays a very important role in evaluation of malignant bowel neoplasms [8].

The objectives of the study were to evaluate the various CT findings in malignant bowel neoplasms, extra-intestinal involvement to stage bowel neoplasms and to accurately characterise the tumour histologically as far as possible.

MATERIALS AND METHODS

A retrospective descriptive study was carried out in the Department of Radiology, Maharajah's Institute of Medical Sciences (MIMS), Nellimarla, Vizianagaram district.

Inclusion criteria: This study included a total of 38 patients of proven bowel malignancy on biopsy who underwent CT examination of the abdomen from November 2014 to May 2017. All those patients who presented with history of melena, constipation, diarrhoea, alternate diarrhoea and constipation, weight loss and anorexia were included in the study.

Exclusion criteria: Patients with synchronous or metachronous malignancy of other organs in addition to bowel malignancy, severely decompensated patients, pregnancy and patients with deranged kidney function test.

Formal sample size calculation was not done. All the data of patents fulfilling the inclusion and exclusion criteria along with informed consent, were enrolled in the study.

Procedure

The patient was placed supine on the CT table. The head was immobilised with head rest. Hands were folded above the shoulder around the head. A scout topogram was obtained from domes of diaphragm down to the symphysis pubis. Appropriate positioning with help of laser mark with respect to the field of interest was done.

Scan parameters included 220 mAs (milliampere-second)/140 (kVp) Kilo voltage peak. Pre-contrast contiguous sequential scans with slice thickness of 10 mm were taken from domes of diaphragm down to the level of pubic symphysis followed by oral and postintravenous contrast scans were done for adequate bowel distension and luminal enhancement. History and physical examination of all patients was systematically carried out. All the patients were required to be fasting for at least 6 hours prior to the examination.

Contrast

Oral contrast: Patients were made to drink 300-400 mL of solution prepared with two ampules of 60% iodinated contrast mixed in 1½ L of plain water at 20-30 minutes intervals with 100-200 mL of same solution to be given on CT table for adequate distension of stomach.

Rectal contrast: To opacify the rectum and distal colon, 200-500 mL of rectal contrast is given rectally as a standard procedure. In female married patients, vaginal tampon is inserted in the vagina to delineate anatomic structures in pelvis.

Post-intravenous contrast: A bolus of 100 mL of 76% iodinated contrast was injected at 1.7 to 2 mL/sec via a pressure injector. Scan delay is 46-50 seconds, collimation 5 mm, table feed 5 mm, reconstruction interval 5 mm, phases of acquisition single phase, dual phase, if needed, region scanned from above domes of diaphragm down to the symphysis pubis. Delayed scans were obtained whenever it was deemed necessary. Multiplanar reconstruction and Three Dimensional (3D) volume rendering techniques can be used to better visualise colon anatomy and the location of the suspected mass or abnormality. Repeated imaging can be done with the patient in prone position, if appropriate or with more air or contrast material after administration to distend collapsed bowel parts. A scan time of 20 minutes was observed for each case.

Normal small-bowel wall is thin, measuring between 1 and 2 mm when the lumen is well-distended. However, depending on the degree of luminal distention, the thickness of the usual small-bowel wall varies slightly. As a consequence, multiple standards for the diagnosis of small-bowel wall thickening have been used. The wall is also not seen when the lumen of the small bowel is distended. If the bowel is incompletely collapsed, the wall measures between 2 and 3 mm and is of symmetric thickness. In these cases it is essential to relate the degree of thickness of similarly distended segments to exclude disorders [9]. The usual thickness of the colonic wall differs greatly depending on the degree of distention. When the colon is distended, the wall should measure less than 3 mm, and it is often imperceptible [9]. The actual thickness is often difficult to assess due to fecal matter, fluid, or colonic redundancy. The true thickness will often be revealed prudently by following the colonic wall to a region where the colon is well-distended with gas. Luminal margins are usually smooth being well-outlined by the contrast/air. The outer colonic margin is sharply outlined by surrounding homogenous pericolic fat [9].

After an appropriate bolus of IV contrast material, the regular intestinal wall strengthens. In patients who have been given water as an oral contrast agent, the enhancement is also more readily detected. In these situations, next to the low-attenuation fluid in the lumen, the strengthened bowel wall is well depicted. On the mucosal part of the bowel wall, enhancement is usually greater. It is not appropriate to mistake this change for a disease phase. Identifying that the wall is not thickened and that there is no perienteric inflammation, will allow one to distinguish normal improvement from the disease process [9]. Fat in mesentery has the same attenuation as fat elsewhere in the body. Major arteries and veins are identified as branching structures within mesenteric fat and do exceed 3 mm in diameter. Mesenteric lymph nodes are occasionally observed in mesentery exceeds 4 mm in normal individual.

CT interpretative criteria for malignant bowel neoplasms which were used in the current study were [9]:

- Site involved (small bowel-jejunum and ileum, large bowel-caecum, colon, rectum and anal canal)
- Length of bowel involved (short segment: less than 10 cm, long segment: more than 10 cm)
- Degree of bowel wall thickening (more than 2 cm and less than 2 cm)
- Circumferential/eccentric bowel wall thickening.
- Attenuation and enhancement pattern of involved bowel wall (Homogenous attenuation and heterogenous attenuation).
- Extra-intestinal involvement includes perienteric/surrounding involvement, lymphadenopathy, and solid organ abnormalities/metastasis.

STATISTICAL ANALYSIS

Statistical data was presented in the form of frequency and percentages for malignant lesions of small and large intestine.

RESULTS

The present study included 38 patients of biopsy proven malignant bowel neoplasm with age between 3 and 70 years. Age groups between 40-59 years formed the bulk of cases (47.36%). Male cases (68.5%) were more than female cases (31.5%) as shown in [Table/Fig-1].

Variables	No. of patients	Percentage
Age group (years)		
0-9	3	7.9%
10-19	1	2.6%
20-29	5	13.1%
30-39	7	18.5%
40-49	9	23.68%
50-59	9	23.68%
60-69	4	10.5%
Sex		
Male	26	68.42%
Female	12	31.57%

[Table/Fig-1]: Age and Sex-wise distribution of cases (N=38).

In small bowel, ileum was the commonest site of involvement (13.15%), whereas rectum (26.3%) was commonest site of involvement in the large bowel [Table/Fig-2]. Based on the length of bowel segment involved, cases with short segment involvement of bowel was more common 27 (71.1%) than long segment involvement 11 (28.9%). Circumferential bowel wall thickening is more common (86.8%) than eccentric wall thickening (13.2%) [Table/Fig-3]. Based on the enhancement pattern of cases, homogenous enhancement pattern was seen in 11 (28.9%) number of cases whereas heterogenous enhancement pattern was seen in 27 (71.1%) cases. Lymph nodal involvement was seen in 44% cases and metastasis to

Site involved	Number of patients	Percentage
Jejunum	1	2.6%
Ileum	5	13.1%
Ileocaecal	3	7.8%
Caecum	1	2.6%
Ascending colon+ caecum	2	5.3%
Ascending colon	4	10.5%
Transverse colon	5	13.3%
Descending colon + sigmoid colon	1	2.6%
Sigmoid colon+Rectum	3	7.8%
Rectum	10	26.3%
Rectum +anal canal	3	7.8%

[Table/Fig-2]: Site of GIT involved in the malignant neoplasm.

Preventive studies	Circumferential wall thickening		Mass lesion/eccentric lesion	Total number of patients
	<2 cm	>2 cm		
Number of patients	5	28	5	38
Percentage	13.2%	73.6%	13.2%	100%


[Table/Fig-3]: Distribution of cases based on circumferential wall thickening.

Distribution of cases	Number of patients	Percentage (%)
Extraintestinal involvement		
Local invasion	4	10.5%
lymph node involvement	17	44.7%
Liver metastasis	11	28.9%
Histological type		
Adenocarcinoma	23	60.5%
Lymphoma	9	23.7%
Mucinous adenocarcinoma	4	10.5%
GIST	2	5.2%


[Table/Fig-4]: Distribution of cases based on extra-intestinal involvement and histological type.

Histopath diagnosis	Length of bowel involved		Degree of bowel wall thickening (more than 2 cm and less than 2 cm)		Attenuation and enhancement pattern of involved bowel wall (Homogenous attenuation and heterogenous attenuation)		Extraintestinal involvement includes perienteric/surrounding involvement, lymphadenopathy, and solid organ abnormalities/metastasis	
	Short segment	Long segment	>2 cm	<2 cm	Homogenous	Heterogenous		
Adenocarcinoma (n=23)	20	3	17	6	8	15	Lymphadenopathy	11
							Liver metastasis	2
Lymphoma (n=9)	6	3	9	0	3	6	Abdominal lymphadenopathy	4
							Splenic deposits	1
							Liver deposits	2
							Lymphadenopathy	2
Mucinous adenocarcinoma (n=4)	4	0	4	0	0	4	Ascites liver scalloping	1
							-	-
GIST (n=2)	2	0	2	0	0	2	-	-


[Table/Fig-5]: Computed Tomography (CT) characteristics with each histopath proven neoplasm. Histopath: Histopathological



[Table/Fig-6]: Rectosigmoid adenocarcinoma with pelvic lymphadenopathy: a) NECT (at the level of pelvis) showing eccentric bowel wall thickening of 3.5 cm in rectosigmoid region (blue arrow) with pelvic lymphadenopathy (red arrow); b) IV Contrast enhanced CT showing heterogenous enhancement of rectosigmoid wall thickening (blue arrow) with rim enhancing pelvic lymphadenopathy (red arrow). NECT: Non-enhance computed tomography; CT: Computed tomography



[Table/Fig-8]: Colonic lymphoma: a) CT with oral contrast (at the level of upper abdomen) showing long symmetric concentric wall thickening (blue arrows) of 2 to 3 cm involving the hepatic flexure and transverse colon. b) IV contrast enhanced CT showing mild heterogenous enhancement of wall thickening (blue arrows). IV: Intravenous



[Table/Fig-7]: Rectal adenocarcinoma with perirectal lymphadenopathy: a and b) CT with rectal contrast (at the level of lower pelvis) showing eccentric rectal wall thickening of 2 to 2.5 cm (red arrows) with perirectal lymphadenopathy (green arrows). CT: Computed tomography

liver was also seen in 28.9% cases [Table/Fig-4]. Adenocarcinoma was the commonest histological subtype seen in highest number of patients 23 (60.6%) followed by lymphoma in 23.6% of cases, mucinous adenocarcinoma in 10.5% cases and few cases of about 2 (5.2%) were GIST [Table/Fig-4]. From the [Table/Fig-5-8] study key findings were reported based on CT characteristics with each histopathologically proven neoplasm.

DISCUSSION

In diagnosing malignant neoplasms of the small and large intestine, numerous studies have shown the accuracy of the CT than traditional clinical-radiographic evaluation methods. In our study of 38 patients of the biopsy proven malignant bowel neoplasm, we evaluated the role of CT imaging in characterising the CT features in malignant bowel neoplasm based on involved intestine, length of the diseased intestinal segment, degree of intestinal wall thickening, extra-intestinal involvement, lymph nodal involvement and solid organ metastasis, etc. The commonly affected age groups were of 40-59 years (47.2%) and males (68.5%) were more affected than females. These findings were in line with the study conducted by Ravi Kumar Y et al., which mentioned most affected age groups as 51 to 60 years (23.1%) and majority of the cases were males (63.5%) [10].

Adenocarcinoma is the commonest malignant neoplasm of the colon. The small intestine is uncommon site for adenocarcinoma. Out of 38 patients, 23 were proven to have adenocarcinoma. The rectum was the commonest site of involvement seen in 13 patients (26.3%), and descending colon was affected less and seen only in one case (2.6%). Twenty out of 23 patients (84%) showed short segment involvement i.e., less than 10 cm. Remaining three patients showed long segment involvement i.e., more than 10 cm. Short segment involvement was commonly seen in malignant lesion than long segment which was seen in inflammatory conditions and these finding were in accordance to the Macari M and Balthazar EJ, [9]. Average diseased bowel wall

thickness seen in 17 out of 23 patients was more than 2 cm. This finding was consistent with the study by Macari M et al., which stated that wall thickening greater than 2 cm is usually present in neoplastic conditions, whereas benign conditions result in bowel wall thickening of less than 2 cm [9]. In the present study, 6 patients even though diseased bowel wall thickness was less than 2cm, they were proven to be malignant. Fifteen out of 23 patients showed heterogeneous mixed patterns of enhancement on post-contrast series whereas 8 patients showed homogenous enhancement patterns. These two patterns of enhancement are usually seen in malignant lesion. In these conditions, other factors like degree of bowel wall thickening, circumferential/eccentric bowel wall thickening, length of the involved segment, and associated extra intestinal abnormalities are important in establishing the correct diagnosis. In cases of neoplasm, homogeneous attenuation correlates with size of the tumour. These findings are consistent with various studies [9,11,12,13]. Six out of 13 patients of rectal adenocarcinoma showed peri-rectal and pre-sacral lymph nodal involvement. Any lymph node in presacral and peri-rectal region is considered malignant irrespective of the size, since inflammatory nodes are never seen in these regions. Other lymph nodes that were involved were mesenteric, portal, aortic, aortocaval, iliac and inguinal group. All these nodes were more than 1 cm in short axis diameter, however whether they were malignant or not, could not be accurately mentioned based on CT features. Eight out of 23 patients of adenocarcinoma showed liver metastasis. This finding was similar as study done by Horton KM et al., [11]. CT has an established role in the detection of hepatic metastases in patients with colorectal cancer.

Nine out of 38 patients were proven to be lymphoma on the biopsy, of which majority (4 out of 9) patients showed involvement of the ileum. Most characteristic finding seen in 6 out of 9 patients of lymphoma was the short segment circumferential involvement with aneurysmal dilatation of the bowel. This dilatation was secondary to loss of muscle tone of the intestinal wall caused by extensive invasion of muscle layer and neural plexus. These findings are consistent with the study of Balthazar EJ et al., [14]. Six out of 9 patients of lymphoma showed heterogenous mixed enhancement pattern and 3 patients showed homogenous pattern. Four out of 9 patients of lymphoma showed enlarged abdominal lymph nodes, two patients showed liver deposits and one patient had splenic deposit.

Four out of 38 patients (10.5%) were proven to be mucinous adenocarcinoma of the large bowel and two out of four patients showed foci of calcifications within mass, two patients showed lymph nodal enlargement one of which showed calcification. One patient showed ascites with scalloping of liver surface. Heterogenous mixed pattern of enhancement was seen in all four patients. Presence of tumoural calcification and calcified lymph nodes along with heterogeneity on post-contrast is common finding in mucinous adenocarcinoma. These findings correlated with the findings by Hussain SM et al., [15]. All the four patients showed short segment involvement.

Two out of 38 patients (5.2%) were proven to be GISTs, one involving the jejunum and another involving the transverse colon. Both these

show varying degree of internal necrosis or cystic changes. Pattern of growth in jejunal GIST was of endoenteric type and that involving the transverse colon was of exocolic type with large exophytic component showing cystic changes.

Limitation(s)

Small sample size, single slice CT machine are the limitations of this study.

CONCLUSION(S)

Computed tomography imaging findings like involved intestine, length of diseased intestinal segment, degree of intestinal wall thickening, extra intestinal involvement with respect to involvement of surrounding structures, lymph nodal involvement, distant metastasis along with histological type in most cases could be found, thus permitting tumour characterisation which can provide very useful guide in diagnosis of malignant neoplasms of small and large intestine. The role of CT in diagnosing and altering the treatment plans for a wide range of causes of small and large intestine neoplasms was assertively determined in this study.

REFERENCES

- [1] Freeman AH. CT and bowel diseases. *BJR*. 2001;74(877):04-14.
- [2] Evers BM. Sabiston textbook of surgery. 18th edition. Noida: Elsevier. 2009; 1289.
- [3] Frager D. Intestinal obstruction role of CT. *Gastro enteral Clin North Am*. 2002;31:777-99.
- [4] Sarr MG, Bulkley GB, Zuidema GD. Preoperative recognition of intestinal strangulation obstruction: prospective evaluation of diagnostic capability. *Am J Surg*. 1983;145(1):176-82.
- [5] Lorgan JG, DuBrow RA. The computed tomographic appearances and clinical significance of intussusception in adults with malignant neoplasms. *Br J Radiol*. 1990;63:257-63
- [6] Megibow AJ, Balthazar EJ, Cho KC, Medwid SW, Birnbaum BA, Noz ME. Bowel obstruction: evaluation with CT. *Radiology*. 1991;180(2):313-18.
- [7] Kim HK, Cho MJ, Ha S, Seo MJ, Moon SG, Oh JH. Adenocarcinoma of the proximal jejunum misdiagnosed as gastroparesis. *The Korean Journal of Helicobacter and Upper Gastrointestinal Research*. 2019;19(3):202-06.
- [8] Jahromi AH, Kim RH, Chu QD, Li BD. Small Bowel Cancer (excluding gastrointestinal stromal tumours and carcinoid): clinical management. *The American Cancer Society's Oncology in Practice*. Wiley publications. USA. 2018; 69-81.
- [9] Macari M, Balthazar EJ. CT of bowel wall thickening: significance and pitfalls of interpretation. *AJR Am J Roentgenol*. 2001;176:1105-16.
- [10] Yeli RK, Patil SV, Nagababu P. Multidetector computed tomography in small and large bowel lesions. *International Journal of Contemporary Medicine Surgery and Radiology*. 2018;3(1):151-54.
- [11] Horton KM, Juluru K, Montgomery E, Fishman EK. Computed tomography imaging of gastrointestinal stromal tumours with pathology correlation. *J Comput Assist Tomogr*. 2004;28(6):811-17.
- [12] Buckley JA, Fishman EK. CT evaluation of small bowel neoplasms: spectrum of disease. *Radiographics*. 1998;18(2):379-92.
- [13] Freeny PC, Marks WM, Ryan JA, Bolen JW. Colorectal carcinoma evaluation with CT: preoperative staging and detection of postoperative recurrence. *Radiology*. 1986;158(2):347-53.
- [14] Balthazar EJ, Noordhoorn M, Megibow AJ, Gordon RB. CT of small-bowel lymphoma in immunocompetent patients and patients with AIDS: comparison of findings. *Am J Roentgenol*. 1997;168(3):675-80.
- [15] Hussain SM, Outwater EK, Siegelman ES. Mucinous versus nonmucinous rectal carcinomas: differentiation with MR imaging. *Radiology*. 1999;213(1):79-85.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Radiology, Maharajah's Institute of Medical Sciences (MIMS), Vizianagaram, Andhra Pradesh, India.
2. Associate Professor, Department of Radiodiagnosis, GITAM Institute of Medical Sciences and Research, GITAM Deemed to be University, Visakhapatnam, Andhra Pradesh, India.
3. Specialist, Department of Radiology, Burjeel Day Surgery Centre, UAE, Abu Dhabi, UAE, United Arab, Emirates.
4. Third Year Resident, Department of Endocrinology, JIPMER, Puducherry, Tamil Nadu, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Devara Anil Kashi Vishnuvardhan,
Assistant Professor, Department of Radiology, Maharajah's Institute of Medical Sciences (MIMS), Nellimarla, Vizianagaram district, Vizianagaram, Andhra Pradesh, India.
E-mail: rajaniagis@gmail.com; anilkashi@yahoo.com

PLAGIARISM CHECKING METHODS: (Lain H et al.)

- Plagiarism X-checker: Sep 25, 2020
- Manual Googling: Feb 04, 2021
- iThenticate Software: Mar 13, 2021 (13%)

ETYMOLOGY: Author Origin

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? No
- Was informed consent obtained from the subjects involved in the study? NA
- For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: **Sep 24, 2020**
Date of Peer Review: **Nov 27, 2020**
Date of Acceptance: **Feb 11, 2021**
Date of Publishing: **Jul 01, 2021**