

Estimation of Stature among Healthcare Workers using Arm Span, Biacromial Breadth, and Forearm Length Dimensions: A Cross-sectional Study

RICHA BHATNAGAR¹, GARIMA SHARMA², ROHIT SHARMA³, VISHAL KALIA⁴,
VIPUL PARMAR⁵, MANOJ KUMAR GANDHI⁶



ABSTRACT

Introduction: Stature is one of the most commonly used anthropometric dimensions for identification purposes in forensic investigations and for assessing the development and growth of individuals. It is well-known that various anatomical anthropometric parameters can accurately estimate stature.

Aim: To estimate stature among healthcare workers in Himachal Pradesh using linear regression equations based on percutaneous anthropometric parameters of the upper limbs and trunk.

Materials and Methods: This cross-sectional study was conducted on healthcare workers at Dr. Rajendra Prasad Government Medical College, Kangra, Tanda, Himachal Pradesh for a one year (September 2021-August 2022) duration. The study included 360 healthy healthcare workers from the Himachal Pradesh population, 180 females and 180 males, aged 21 years and above. Stature, arm span, Biacromial Breadth (BAB), and

forearm length dimensions were measured for each subject. Statistical analysis was done using Microsoft Excel and Epi-Info version 7.1 software. The Pearson Correlation Coefficient was used to determine the correlation between stature and different anatomical anthropometric parameters. Linear regression equations were developed to estimate stature, with significance set at $p\text{-value} \leq 0.001$.

Results: The mean age of the study population was 41.33 ± 28.91 years. The descriptive anthropometric parameters measured in the study were as follows: stature 162.4 ± 6.85 and 149.4 ± 7.15 , arm span 165.67 ± 7.5 and 150.31 ± 8.33 , right forearm length 26.68 ± 1.33 and 24.36 ± 1.59 , left forearm length 26.47 ± 1.30 and 24.30 ± 1.55 , BAB 34.11 ± 2.16 and 31.74 ± 2.32 , respectively for males and females.

Conclusion: The regression equations derived from this study can be valuable for estimating stature in situations where accurate stature measurements are not feasible.

Keywords: Anthropometry, Anthropometric parameters, Regression, Upper limb

INTRODUCTION

Identification is an individual's birth-right. Anthropometry, which deals with measurements of humans, has been widely used in many clinical and forensic investigations. It provides us with the scientific basis for estimating various measurements in living and deceased individuals [1]. Of all the anthropometric dimensions, stature, measured as the standing height of the individual, is the most commonly used [2]. Apart from aiding in identification, stature measurement is essential for therapeutic purposes such as estimating nutritional status indicators, assessing children's growth, and adjusting drug dosages [3]. However, in some cases where patients are unable to stand, individuals have trunk or leg deformities or lower limb amputations, patients with contractures or fractures, and in mutilated bodies, measuring stature is difficult or impossible [4].

Although the anatomical method is the most precise method for stature estimation, it cannot be used by forensic experts when mutilated bodies or skeletal remains are found in cases of homicide to conceal identity or in cases of mass disasters. In such cases, the mathematical method can be used for stature estimation [5]. Stature reconstruction from the upper limb bones, mainly long bones, is of paramount medico-legal importance. A strong association is seen linking stature with the long bones of the upper extremity [6]. To derive regression formulas for estimating stature, long bones are usually assumed to be more precise than the bones of the foot and hand [7]. Karl Pearson developed the first formal stature regression formula to estimate stature [8]. A method of estimating stature from long bones was also developed by Trotter M and Gleser GC [9].

They derived regression equations from the length of long bones and calculated stature. This is the most common method for stature estimation. Inter-population comparison and estimation of stature from these formulas cannot be generalised as climate, heredity, and the nutritional status of a specific population have been reported to have an effect on stature [10].

Due to various ethnic and racial variations among the Indian population, area-wise, population-specific anthropometric studies are essential in India. The population of Himachal Pradesh is an admixture of two races, Mongoloid and Caucasian. Hence, the data of the country cannot be very reliable for the current population under study. Data for anthropometry of the population of Himachal Pradesh is limited in the literature. Therefore, the present study has attempted to determine stature from arm span, forearm length, and BAB measurements and then devise adequate gender-specific regression analysis for this purpose.

MATERIALS AND METHODS

This cross-sectional study was conducted in the Department of Anatomy at Dr. Rajendra Prasad Government Medical College, Kangra in Tanda, Himachal Pradesh, India for one year from 2021 to 2022, after obtaining ethical committee approval for the study (No. HFW-H DRPGMC/Ethics/2021/25). It included 360 healthy healthcare workers from the Himachal Pradesh population, with 180 females and 180 males, aged above 21 years.

Sample size calculation: Participants were selected using simple random sampling method. In a study published by Aggarwal AN et

al., among North Indian subjects, the mean height and arm span ratio among males and females was 0.9819 ± 0.0239 and 0.9711 [11]. With the above values and taking α as 0.05 and 95% power a sample size of 282 (141 in each group) was obtained.

Inclusion criteria: Only healthy healthcare workers belonging to both male and female genders, of Himachal Pradesh population with age above 21 years, free from any developmental defects or skeletal abnormalities were included in the study.

Exclusion criteria: Healthcare workers with any orthopaedic deformity like bone fracture, hormonal disorders like gigantism and acromegaly, genetic disorders like noonan syndrome, osteogenesis imperfecta or any other disorders that could directly affect the measuring part of the current study and stature of an individual were excluded.

Study Procedure

Study participants were explained the study purpose in their local language and informed consent was taken.

Calibrated Stadiometer (SDM210) (Lifeyelid enterprises) with length of 20 cm to 205 cm (graduation.1cm) was used [Table/Fig-1] to measure stature in centimeters. Stature was measured from vertex to the floor in standing position with the subject standing bare foot on the baseboard of stadiometer and Frankfurt's plane parallel to the ground [12].



[Table/Fig-1]: Stature measurement.


Arm span was measured [Table/Fig-2] in centimeters as the distance from the tip of the middle finger of one arm to the tip of the middle finger of the other arm, with the subject standing erect against the wall and arms outstretched, palms facing forwards [13].

Forearm length was measured [Table/Fig-3] in centimeters as the distance between the tip of the styloid process of the ulna and the tip of the olecranon process in a fully flexed elbow, with the palm facing over the opposite shoulder [14]. BAB was measured [Table/Fig-4] in centimeters by first feeling the outside edge of the acromial process and then measuring the distance between the tips of the right and left acromion processes of the shoulders [15].


Arm span, forearm length, and BAB measurements were taken using an Anthropometer with a length of 0-210 cm (BAI 101 Biotech and Scientific Instruments). All measurements were taken thrice, and the mean was calculated for each reading to avoid errors. To minimise errors due to stature and daytime variations, the measurements were taken by one person using the same measuring instruments before noon at a specified fixed time [16-18].



[Table/Fig-2]: Arm span measurement.



[Table/Fig-3]: Forearm length measurement.



[Table/Fig-4]: Biacromial Breadth (BAB) measurement.

STATISTICAL ANALYSIS

Microsoft Excel and Epi-Info version 7.1 software were used for statistical analysis. Mean and standard deviation were used to present quantitative variables, while qualitative variables were presented in the form of percentage and frequency. Pearson's correlation coefficient was calculated to determine the correlation between stature and arm

span, forearm length, and BAB. The coefficient of determination (R^2) was calculated to determine the strength of association among the parameters considered [19]. Gender-specific regression equations, as well as separate ones for combined cases, were developed by a linear regression model to predict stature from an individual's arm span, forearm length, and BAB.

RESULTS

The observations were conducted on 180 males and 180 females, totaling 360 subjects [Table/Fig-5]. All the studied anthropometric parameters, including stature, arm span, forearm length (right and left sides), and BAB, were found to be higher in male subjects compared to females (p -value<0.001) [Table/Fig-6]. Pearson's correlation coefficient (r) indicated a positive correlation (p -value<0.001) between stature and the parameters studied [Table/Fig-7]. Higher R^2 (coefficient of determination) values indicated that stature can be accurately predicted from arm span, forearm length, and BAB in decreasing order. Left forearm measurements significantly predicted stature compared to right forearm.

Gender	Mean age (in years)	SD
Males (n=180)	41.07	10.3
Females (n=180)	41.58	7.34
Total	41.33	8.91

[Table/Fig-5]: Distribution of cases according to their age (n=360).

Variable (in centimeters)	Male (N=180)		Female (N=180)		Combined (N=360)		*p-value
	Range	Mean±SD	Range	Mean±SD	Range	Mean±SD	
Stature	148-176	162.4±6.85	136-168	149.4±7.15	136-176	155.9±9.56	<0.001
Arm span	151-181	165.67±7.5	133-168.5	150.31±8.33	133-181	157.99±11.04	<0.001
Right forearm length	24.7-30	26.68±1.33	21.1-27.7	24.36±1.59	21.1-30	25.52±1.87	<0.001
Left forearm length	24.3-29.9	26.47±1.3	21.2-27.9	24.3±1.55	21.2-29.9	25.38±1.8	<0.001
Biacromial breadth (BAB)	28.5-38.7	34.11±2.16	26.2-36.4	31.74±2.32	26.2-38.7	32.93±2.53	<0.001

[Table/Fig-6]: Descriptive statistics of anthropometric parameters measurements in study subjects.

#p-value between male and female; Independent student t-test of significance for comparing means and standard deviation; N: number of subjects; SD: Standard deviation

Following regression analysis, equations were derived for estimating stature from arm span, forearm length, and BAB for males, females, and combined cases [Table/Fig-7]. Scatter plots illustrating the

Variable	r	R ²	Regression equation Y=A+BX	p-value
Arm span (AS)				
Male	0.861	0.741	Stature= 32.39+0.78×AS	<0.001
Female	0.897	0.804	Stature= 33.64+0.77×AS	<0.001
Combined	0.937	0.878	Stature= 27.72+0.81×AS	<0.001
Right forearm length (RFAL)				
Male	0.7	0.490	Stature= 66.79+3.58×RFAL	<0.001
Female	0.711	0.506	Stature= 71.89+3.18×RFAL	<0.001
Combined	0.828	0.685	Stature= 48.08+4.22×RFAL	<0.001
Left forearm length (LFAL)				
Male	0.732	0.536	Stature= 59.52+3.88×LFAL	<0.001
Female	0.752	0.565	Stature= 65.45+3.45×LFAL	<0.001
Combined	0.845	0.714	Stature= 41.51+4.5×LFAL	<0.001
Biacromial Breadth (BAB)				
Male	0.542	0.294	Stature =103.97+1.71×BAB	<0.001
Female	0.458	0.209	Stature =104.68+1.4×BAB	<0.001
Combined	0.642	0.412	Stature =76.19+2.42×BAB	<0.001

[Table/Fig-7]: Regression equation for estimating stature from anatomical anthropometric parameters in study subjects.

r: Correlation coefficient, R^2 : Coefficient of determination; Y: stature in cm (dependent variable); A: constant; B: Regression coefficient of variable (linear regression model) and X: independent variable (arm span, forearm length, BAB in cm); AS: Arm span; RFAL: Right forearm length; LFAL: Left forearm length; BAB: Bi Acromial breadth.


linear correlation of stature and arm span in males and females, stature and forearm length in males, and stature and forearm length in females are presented in [Table/Fig-8-10].

The accuracy of measured stature was assessed by comparing the mean value of measured stature using the regression formula with the mean value of the actual stature of the study subjects. The similarity of their values demonstrated that the regression equations formulated in the present study could be utilised as a reliable tool for estimating stature from arm span, forearm length, and BAB measurements. For example, in a female with measured parameters as follows: arm span= 163 cm, right forearm length= 27.5 cm, left forearm length= 27.4 cm, and BAB= 35.2 cm, the estimated stature obtained using these measurements in the regression equations was 159.2 cm, 159.3 cm, 159.9 cm, and 154 cm, respectively. The actual stature of that female was 160 cm, confirming that the derived equations can be accurately used for stature estimation.


[Table/Fig-11] showed scatter plot diagrams depicting linear correlation of stature and Biacromial Breadth (BAB) in males and females.

DISCUSSION


All parts of the body exhibits biological correlation, where the size of one part can be used to estimate the size of another. Similarly, stature can be accurately estimated from various body parts. Several factors, including age, sex, and race, influence skeletal development and various body measurements [20,21]. Stature estimation is




[Table/Fig-8]: Scatter plots depicting linear correlation of stature and arm span in males and females.



[Table/Fig-9]: Scatter plots depicting linear correlation of stature and forearm length in males.



[Table/Fig-10]: Scatter plots depicting linear correlation of stature and forearm length in females.



[Table/Fig-11]: Scatter plots depicting linear correlation of stature and Biacromial Breadth (BAB) in males and females.

crucial for identification purposes in anthropological studies and medico-legal examinations. Stature estimation can be performed accurately even in fragmentary remains and in dismembered and mutilated bodies [10]. Various researchers have attempted to estimate stature from different body parts.

Anthropometric measurements in males were found to be greater than in females, which can be attributed to the genetic make-up of males. The pubertal age of males is typically two years later than that of females, providing them with additional growth opportunities [10]. The same regression equations cannot be applied to both male and female cases due to the higher mean values of parameters observed in male subjects compared to females (p -value <0.001).

The correlation coefficient (r) for stature and arm span was 0.861 in males and 0.897 in females, which was statistically highly significant (p -value <0.001) and indicated a strong correlation. Similar findings have been reported in other studies conducted globally [Table/Fig-12] [10,22-25]. A similar study on estimation of stature from outstretched arm span and measurement of components of upper limb in the natives of Gujarat [26] to further support the findings.

While some researchers, like Potdar AB et al., only took one-sided forearm measurements for stature correlation [27], the present study involved measurements from both sides for improved stature correlation. Although a significant correlation with stature was observed in both right and left forearm measurements, the left forearm measurement showed a more significant correlation with stature compared to the right. A comparison with different studies on forearm measurements is depicted in [Table/Fig-13] [28,29].

A review of the literature revealed that very few researchers have attempted to correlate BAB with stature [Table/Fig-14] [15,30-32]. In current study, a higher correlation coefficient (r) was observed in male cases compared to females. This difference could be explained by the fact that in males, the shoulder growth spurt at the beginning of puberty is more distinct, leading them to develop broader shoulders compared to females [30].

Study	Sex	r	R ²	Regression Equation
Supare MS et al., [10] (2015)	Male	0.89	0.79	Stature=31.26+0.806 × AS (cm)
	Female	0.90	0.81	Stature=33.54+0.77 × AS (cm)
Uzun Ö et al., [22] (2018)	Male	0.785	0.716	Stature=420.527+ 0.755×AS (cm)
	Female	0.807	0.751	Stature=432.536+ 0.736×AS (cm)
Mumtaz SH et al., [23] (2019)	Male	0.766	-	Stature= 33.340 + 0.784×AS (cm)
	Female	0.887	-	Stature= 20.796+1.109×AS (cm)
Sarma A et al., [24], (2020)	Male	0.988	0.976	Stature=1.060+0.954×AS (cm)
	Female	0.991	0.983	Stature=0.150+0.962×AS (cm)
Rai P et al., [25] (2021)	Male	0.781	0.610	Stature= 40.75+0.7348×AS (cm)
	Female	0.837	0.700	Stature= 47.001+ 0.6712×AS (cm)
Present study	Male	0.861	0.741	Stature= 32.39+0.78 × AS (cm)
	Female	0.897	0.804	Stature= 33.64+0.77 × AS (cm)

[Table/Fig-12]: Comparative data of different studies on stature estimation from arm span [10,22-25].

Study	Sex	Forearm length LFAL/RFAL	r	R ²	Regression equation
Londhe S et al., [28] (2019)	Male	RFAL	0.514	-	Stature=95.82+2.887×RFAL (cm)
		LFAL	0.501	-	Stature=116.5+2.128×LFAL (cm)
	Female	RFAL	0.598	-	Stature=99.46+2.427×RFAL (cm)
		LFAL	0.615	-	Stature=95.08+2.632×LFAL (cm)
Dongre SS et al., [29] (2021)	Male	RFAL	0.969	0.941	Stature=126.42+1.4296×RFAL (cm)
		LFAL	0.979	0.959	Stature=128.32+1.3817×LFAL (cm)
	Female	RFAL	0.974	0.949	Stature=127.26+1.3852×RFAL (cm)
		LFAL	0.984	0.967	Stature=128.52+1.3711×LFAL (cm)
Present study	Male	RFAL	0.7	0.490	Stature=66.79+3.58×RFAL (cm)
		LFAL	0.732	0.536	Stature=59.52+3.88×LFAL (cm)
	Female	RFAL	0.711	0.506	Stature=71.89+3.18×RFAL (cm)
		LFAL	0.752	0.565	Stature=65.45+3.45×LFAL (cm)

[Table/Fig-13]: Comparative data of different studies on stature estimation from forearm length [28,29, Present study].

Study	Sex	r	R ²	Regression equation
Ozaslan A et al., [15] (2011)	Male	0.42	0.17	Stature=125.11+1.226×BAB (cm)
	Female	0.26	0.07	Stature=133.16+0.822×BAB (cm)
Patel SH et al., [32] (2015)	Male	0.31	-	Stature=140.01+0.71×BAB (cm)
	Female	0.23	-	Stature=137.31+0.50×BAB (cm)
Parul et al., [30] (2019)	Male	0.492	-	Stature=122.37+1.389×BAB (cm)
	Female	0.313	-	Stature=130.84+0.746×BAB (cm)
Yadav S et al., [31] (2023)	Male	0.369	-	Stature=45.72+3.108×BAB (cm)
	Female	0.206	-	Stature=135.16+0.549×BAB (cm)
Present study	Male	0.542	0.294	Stature=103.97+1.71×BAB (cm)
	Female	0.458	0.209	Stature=104.68+1.4×BAB (cm)

[Table/Fig-14]: Comparative data of different studies on stature estimation from Biacromial Breadth (BAB) [15,30-32].

Various factors such as nutritional status, sex, and age influence stature and body dimensions. Additionally, genetic, environmental, and racial factors play a role in influencing an individual's skeletal growth [33].

Limitation(s)

The derived equations are specific to the population studied, and if applied to other populations, they may not yield accurate results. Furthermore, the equations will provide accurate results only when applied to individuals older than 21 years. Additionally, the equations have limited applicability in individuals with any disorder or defect that affects skeletal growth.

CONCLUSION(S)

Among all the parameters studied, stature is most accurately determined by arm span, followed by forearm length, and least by

BAB in both sexes. The gender-specific and combined regression equations derived in the study to estimate stature from arm span, forearm length, and BAB can be effectively used in individuals for whom precise measurement of stature is not feasible. This will assist in identification and can be beneficial to anthropologists, medico-legal experts, and anatomists.

REFERENCES

- [1] Patel SM, Doshi V, Ruparelia S, Ankur Z, Dhara P, Rathod SP, et al. Anthropological study of the foot and it's relationship between different parameters and stature in an adult population of different areas of Gujarat. *Natl J Integr Res Med*. 2011;2(3):62-65.
- [2] Heymsfield SB, Gallagher D, Mayer L, Beetsch J, Pietrobelli A. Scaling of human body composition to stature: New insights into body mass index. *Am J Clin Nutr*. 2007;86(1):82-91.
- [3] Kaestner SA, Sewell GJ. Chemotherapy dosing part I: Scientific basis for current practice and use of body surface area. *J Clin Oncol*. 2007;19(1):23-37.
- [4] Chawla M, Rajkumar, Tomar S, Ashoka R. The relationship between arm span and height in adult males of north Indian Punjabi population. *J Evol Med Dent Sci*. 2013;2(4):332-39.
- [5] Duyar I, Can P, Zagypar R. A new method of stature estimation for forensic anthropological application. *Athnthropological Science*. 2006;114(1):23-27.
- [6] Nandi ME, Olabiyo O, Ibeabuchi NM, Okubike AE, Cyprian EI. Stature reconstruction from percutaneous anthropometry of long bones of upper extremity of Nigerians in the university of Lagos. *Arab Journal of Forensic Sciences and Forensic Medicine*. 2018;1(7):869-80.
- [7] Singh S, Nair SK, Anjankar V, Bankawar V, Satpathy DK, Malik Y. Regression equation for estimation of femur length in central Indians from inter-trochanteric crest. *Journal of Indian Academy of Forensic Medicine*. 2013;35(3):223-26.
- [8] Karl P. Mathematical contributions to the theory of evolution. On the reconstruction of the stature of prehistoric races. *Philos Trans A Math Phys Eng Sci*. 1898;192:169-244. 10.1098/rsta.1899.0004
- [9] Trotter M, Gleser GC. A re-evaluation of estimate of stature based on measurements of stature taken during life and of long bones after death. *Am J Phys Anthropol*. 1958;16(1):79-123.
- [10] Supare MS, Bagul AS, Pandit SV, Jadhav JS. Estimation of stature from arm span in medical students of Maharashtra, India. *Ann Med Health Sci Res*. 2015;5(3):218-21.
- [11] Aggarwal AN, Gupta D, Ezekiel LM, Jindal SK. Statistical estimation of height from arm span in north Indian subjects. *Indian J Physiol Pharmacol*. 2000;44(3):329-34.
- [12] Uhrová P, Beňuš R, Masnicová S. Stature estimation from various foot dimensions among Slovak population. *J Forensic Sci*. 2013;58(2):448-51.
- [13] Mulu A, Sisay B. Estimation of stature from arm span, arm length and tibial length among adolescents of aged 15-18 in Addis Ababa, Ethiopia. *Ethiop J Health Sci*. 2021;31(5):1053-60.
- [14] Yeasmin N, Hossain I, Chowdhury MA, Rocky MH, Rouf A, Sokharanjan S. Stature estimation using ulnar length and shoulder elbow length in a Bangladeshi population. *SN Comprehensive Clinical Medicine*. 2020;2(12):2754-62.
- [15] Ozaslan A, Karadayi B, Kulusayin MO, Kaya A. Stature estimation from bi-acromial and bi-iliocristal measurements. *Rom J Leg Med*. 2011;19(3):171-76.
- [16] Singh S, Hussain M, Rizvi SJ. Estimation of stature from foot measurements in adults. *Int J Med Toxicol Leg Med*. 2010;13(2):46-49.
- [17] Rastogi P. Methods of stature estimation: A review. *Indian J Forensic Med Toxicol*. 2009;26(1):10-13.
- [18] Bal K, Bapuli C. Height estimation of an individual from forearm length in medical students. *Asian J Med Sci*. 2022;13(5):89-93.
- [19] Ozaslan A, İçcan MY, Ozaslan I, Tuğcu H, Koç S. Estimation of stature from body parts. *Forensic Sci Int*. 2003;132(1):40-45.
- [20] Methapatil SS, Dethe VA. Prediction of height using hand and foot parameters in Maharashtra population. *Natl J Clin Anat*. 2022;11(3):143-47.
- [21] Krishan K. Anthropometry in forensic medicine and forensic science- 'Forensic anthropometry'. *The Internet Journal of Forensic Science*. 2007;2(1):01-17.
- [22] Uzun Ö, Yeginoglu G, Öksüz CE, Kalkışım ŞN, Zihni NB. Stature estimation utilising arm span measurements in Turkish adults. *J Clin Diagn Res*. 2018;12(12):06-10.
- [23] Mumtaz SH, Quadri SM, Mirza M, Hamid S. Arm span as a predictor of stature. *Int J Sci Res (Ahmedabad)*. 2019;8(6):37-39.
- [24] Sarma A, Barman B, Das GC, Saikia H, Momin AD. Correlation between the arm-span and the standing height among males and females of the Khasi tribal population of Meghalaya state of North-Eastern India. *J Family Med Prim Care*. 2020;9(12):6125-29.
- [25] Rai P, Das A, Kumar S, Girdhar R, Kumar P, Saini N. Is arm-span an accurate measure of stature? A cross sectional study in North Indian population. *Asian Journal of Medical Sciences*. 2021;12(7):112-16.
- [26] Lalhminghluia R, Saiyed MZG, Nayak M, Jani CB. Estimation of stature from outstretched arm span and measurement of components of upper limb in the natives of Gujarat. *Indian Journal of Forensic and Community Medicine*. 2021;8(2):84-90.
- [27] Potdar A, Rathod PI, Potdar PA, Desai MM. A study of estimation of stature from forearm length. *Indian J Forensic Med Toxicol*. 2019;13(2):219-21.
- [28] Londhe S, Panjakash S, Mirzanaiik AD. Stature estimation from forearm lengths in North Karnataka population; India. *Indian Journal of Clinical Anatomy and Physiology*. 2019;6(1):32-37.
- [29] Dongre SS, Deokar RB, Patil SS. Correlation of the stature to forearm length in the young adults of western Indian population. *Medico Legal Update*. 2021;21(1):1062-69.
- [30] Parul, Jain M, Shukla L, Kaur D. Biacromial breadth: A tool to measure stature. *Journal of Medical Science and Clinical Research*. 2019;7(8):644-48.
- [31] Yadav S, Nayak VC, Palimar V, Gupta C, Chetal JR. An estimation of stature and sex using various anthropometric parameters in south Indian population. *Journal of Forensic Science and Medicine*. 2023;9(2):130-36.
- [32] Patel S, Bastia BK, Kumar L, Senthil K. Estimation of adult human stature from measurement of inter-acromial length in Gujarati population of India. *Journal of Indian Academic of Forensic Medicine*. 2015;37(4):365-68.
- [33] Weaver CM, Fuchs RK. Skeletal growth and development. In: Burr DB, Allen MR, editors. *Basic and applied bone Biology*. Philadelphia: Elsevier; 2014. Pp. 245-60.

PARTICULARS OF CONTRIBUTORS:

1. Postgraduate Student, Department of Anatomy, DRRPGMC, Kangra, Himachal Pradesh, India.
2. Assistant Professor, Department of Anatomy, DRRPGMC, Kangra, Himachal Pradesh, India.
3. Assistant Professor, Department of Anatomy, DRRPGMC, Kangra, Himachal Pradesh, India.
4. Professor, Department of Anatomy, DRRPGMC, Kangra, Himachal Pradesh, India.
5. Senior Resident, Department of Forensic Medicine, Dr Radhakrishnan Government Medical College, Himachal Pradesh, India.
6. Senior Resident, Department of Community Medicine, DRRPGMC, Kangra, Himachal Pradesh, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr Vishal Kalia,
Department of Anatomy, DRRPGMC Tanda, Kangra, Himachal Pradesh-176002,
India.
E-mail: drvishalkalia29@yahoo.com

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Jul 21, 2023
- Manual Googling: Jan 31, 2024
- iThenticate Software: Feb 07, 2024 (9%)

ETYMOLOGY: Author Origin

EMENDATIONS: 6

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Jul 21, 2023**

Date of Peer Review: **Oct 24, 2023**

Date of Acceptance: **Feb 08, 2024**

Date of Publishing: **Mar 01, 2024**